



Patient Information – Post-operative care

Mastectomy

Wound care

Your breast tissue has been removed during surgery. The incision has been temporarily numbed with local anaesthetic and sewn together with dissolvable stitches under the skin.

An adhesive dressing has been applied to the wound. You can safely get the dressing wet in the shower and gently dab dry afterward.

A tube has been inserted to drain mastectomy wound fluid. After the drain tube has been removed, typically five to 10 days following surgery, leakage of fluid might occur for a few days through the skin hole. Do not be alarmed. Simply keep the skin clean (wash the area with water) and re-apply an absorptive dressing over the hole.

Please check the wound dressing daily. If you go home with a drain tube in, clear instructions will be given to you by the nursing staff and/or breast care nurse.

Pain relief

Regular paracetamol (e.g. Panadol®), 1g (two 500mg tablets), four times a day, for the first few days after leaving hospital and prior to exercises or functional activities, provides sufficient pain relief for most people. It is safe to add an anti-inflammatory drug (e.g., Nurofen®) for most patients, but please check with your doctor if you have any concerns.

Activity

The physiotherapist will see you on the ward and commence exercises to aid recovery. Gentle exercise, for example walking, is recommended to reduce the risk of deep vein thrombosis. You should avoid more vigorous exercise for at least two weeks after surgery. Driving should be avoided if possible until review following

discharge. You should anticipate at least two weeks off work, but naturally this can vary and will depend on other pending treatments.

Complications

Internal wound bleeding (haematoma) may be suspected if your wound becomes painful, tense, bruised or swollen. Soft bruising alone is of no great concern.

Infection may be suspected if the wound becomes increasingly tender & inflamed. You might start to feel unwell with a fever. This requires urgent attention.

A fluid collection (seroma) is common after lymph node surgery. The chest wall or armpit may become swollen and uncomfortable. Drainage with a needle and syringe is often performed in the doctor's rooms after surgery, to relieve the fluid pressure.

Linear cordes can sometimes be felt along the inside of the arm due to inflamed lymphatics just under the skin. This can be uncomfortable but will improve over the coming weeks or months. It is safe to exercise the arm with lymphatic cording.

Appointment

Your follow-up appointment will be made at the time of your surgery booking.

Please ring the rooms (03 9021 8833) if you do not have an appointment.

Contact

Any concerns, please contact Dr Sarah Birks or the hospital.

Cabrini Brighton

Suite 6, 243 New St,
Brighton 3186

www.drSarahBirks.com.au

Cabrini Malvern Specialist Centre

Level 2, Area E, 183 Wattletree
Rd, Malvern 3144

contact@drsarahbirks.com.au

Waverley Breast Care

357 Blackburn Rd,
Mt Waverley 3149

Ph: 03 9116 4770 **F:** 03 9116 4777